

IIT-GATE: 2011

Chemistry

❖ Question Paper

Section-A

Q.1 - Q.25 carry one mark each.

Q.1 Jahn–Teller distortion of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ acts to

- (a) Raise symmetry.
- (b) Remove an electronic degeneracy.
- (c) Cause loss of H_2O ligand.
- (d) Promote a d–electron to an antibonding molecular orbital.

Q.2 Among the following, the group of molecules that undergoes rapid hydrolysis is

- (a) SF_6 , Al_2Cl_6 , SiMe_4
- (b) BCl_3 , SF_6 , SiCl_4
- (c) BCl_3 , SiCl_4 , PCl_5
- (d) SF_6 , Al_2Cl_6 , SiCl_4

Q.3 The reaction of solid XeF_2 with AsF_5 in 1:1 ratio affords

- (a) XeF_4 and AsF_3
- (b) XeF_6 and AsF_3
- (c) $[\text{XeF}]^+ [\text{AsF}_6]^-$
- (d) $[\text{Xe}_2\text{F}_3]^+ [\text{AsF}_6]^-$

Q.4 A well-known naturally occurring organometallic compound is

- (a) Vitamin B 12 coenzyme
- (b) Chlorophyll
- (c) Cytochrome P–450
- (d) Myoglobin

Q.5 The complex that exists as a pair of enantiomers is

- (a) $\text{trans-}[\text{Co}(\text{H}_2\text{NCH}_2\text{CH}_2\text{NH}_2)_2\text{Cl}_2]^+$
- (b) $\text{cis-}[\text{Co}(\text{NH}_3)_4\text{Cl}_2]^+$
- (c) $[\text{Pt}(\text{PPh}_3)(\text{Cl})(\text{Br})(\text{CH}_3)]^-$
- (d) $[\text{Co}(\text{H}_2\text{NCH}_2\text{CH}_2\text{NH}_2)_3]^{3+}$


Q.6 The region of electromagnetic spectrum employed in the electron spin resonance (ESR) spectroscopy is

- (a) Radiowave (b) Microwave (c) Infrared (d) Visible

Q.7 The red color of oxyhaemoglobin is mainly due to the


- (a) $d-d$ transition (b) Metal to ligand charge transfer transition
(c) Ligand to metal charge transfer transition (d) intraligand $\pi-\pi^*$ transition

Q.8 The band structure in an n-type semiconductor is







- (a) I (b) II (c) III (d) IV


Q.9 In the following reaction




the major product [X] is

- (a)  (b)  (c)  (d) 


Q.10 In the following reaction sequence



the major product [X] is




Q.11 The diene which undergoes Diels-Alder reaction with maleic anhydride is



Q.12 The sequence of an mRNA molecule produced from a DNA template strand with the composition 5'-AGCTACACT-3' is

- (a) 5'-AGUGUAGCU-3' (b) 5'-UCGAUGUGA-3'
- (c) 5'-AGTGTAGCT-3' (d) 5'-TCGATGTGA-3'

Q.13 In the following reaction




the major product [X] is

- (a) (b)
- (c) (d)


Q.14 The structure of the dipeptide Ala-Pro derived from the natural amino acids is

- (a) (b)
- (c) (d)


Q.15 In the following reaction




the major product [X] is



Q.16 In the following reaction



the major product [X] is



Q.17 For a given first order reaction, the reactant reduces to $1/4^{\text{th}}$ its initial value in 10 minutes. The rate constant of the reaction is

- (a) 0.1386 min^{-1} (b) 0.0693 min^{-1} (c) $0.1386 \text{ mol L}^{-1} \text{ min}^{-1}$ (d) $0.0693 \text{ mol L}^{-1} \text{ min}^{-1}$

Q.18 The freezing point constant for water is $1.86 \text{ K (mol kg}^{-1}\text{)}^{-1}$. The change in freezing point when 0.01 mol glucose is added to 1 kg water is

- (a) 1.86 K (b) -1.86 K (c) 0.186 K (d) -0.0186 K

Q.19 On the pressure–temperature diagram for a one–component system, the point where the solid–liquid and the liquid–gas curves intersect is

- (a) Triple point (b) Critical point (c) Melting point (d) Boiling point

Q.20 The wave function for a Harmonic oscillator described by $N x \exp(-\alpha x^2/2)$ has

- (a) One maximum only.
- (b) One maximum, one minimum only.
- (c) Two maxima, one minimum only.
- (d) Two maxima, two minima only.

Q.21 If an arbitrary wave function is used to calculate the energy of a quantum mechanical system, the value calculated is never less than the true energy.

The above statement relates to

- (a) Perturbation theory.
- (b) Variation principle.
- (c) Heisenberg's uncertainty principle.
- (d) Quantization of energy.


Q.22 The point group symmetry of the given planar shape is



Q.23 $\left(\frac{\partial G}{\partial p}\right)_T =$

- (a) V (b) S (c) $-S$ (d) $-V$

Q.24



According to the Ehrenfest classification of phase transitions, the above diagram refers to

- (a) Zeroth order phase transition
- (b) First order phase transition.
- (c) Second order phase transition.
- (d) λ transition.



Q.25 According to conventional transition state theory, for elementary bimolecular reactions, the molar entropy of activation ΔS^{\ddagger} is

- (a) Positive.
- (b) Zero.
- (c) Negative.
- (d) Positive for endothermic and negative for exothermic reactions

Q.26 to Q.55 carry two marks each.

Q.26 The crystal field stabilization energy (CFSE) value, for $[\text{Ti}(\text{H}_2\text{O})_6]^{3+}$ that has an absorption maximum at 492 nm is

- (a) $20,325 \text{ cm}^{-1}$
- (b) $12,195 \text{ cm}^{-1}$
- (c) $10,162 \text{ cm}^{-1}$
- (d) $8,130 \text{ cm}^{-1}$

Q.27 For Et_2AlX ($\text{X} = \text{PPh}_2^-$, Ph^- , Cl^- and H^-), the tendency towards dimeric structure follows the order

- (a) $\text{PPh}_2^- > \text{Cl}^- > \text{H}^- > \text{Ph}^-$ (b) $\text{Cl}^- > \text{PPh}_2^- > \text{H}^- > \text{Ph}^-$
 (c) $\text{Ph}^- > \text{H}^- > \text{Cl}^- > \text{PPh}_2^-$ (d) $\text{H}^- > \text{Ph}^- > \text{PPh}_2^- > \text{Cl}^-$

Q.28 In the isoelectronic series, VO_4^{3-} , CrO_4^{2-} and MnO_4^- , all members have intense charge transfer (CT) transitions. The INCORRECT statement is

- (a) CT transitions are attributed to excitations of electrons from ligand (σ) to metal (e).
 (b) MnO_4^- exhibits charge transfer at shortest wavelength among the three.
 (c) The wavelengths of transitions increase in the order $\text{VO}_4^{3-} < \text{CrO}_4^{2-} < \text{MnO}_4^-$.
 (d) The charge on metal nucleus increases in the order $\text{VO}_4^{3-} < \text{CrO}_4^{2-} < \text{MnO}_4^-$.

Q.29 The increasing order of wavelength of absorption for the complex ions:

- (i) $[\text{Cr}(\text{NH}_3)_6]^{3+}$ (ii) $[\text{CrCl}_6]^{3-}$, (iii) $[\text{Cr}(\text{OH}_2)_6]^{3+}$, (iv) $[\text{Cr}(\text{CN})_6]^{3-}$, is
 (a) $\text{iv} < \text{ii} < \text{i} < \text{iii}$ (b) $\text{iv} < \text{iii} < \text{ii} < \text{i}$ (c) $\text{iv} < \text{i} < \text{iii} < \text{ii}$ (d) $\text{ii} < \text{iii} < \text{i} < \text{iv}$

Q.30 The total number of metal-metal bonds in $\text{Ru}_3(\text{CO})_{12}$ and $\text{Co}_4(\text{CO})_{12}$, respectively, is

- (a) 3 and 6 (b) 4 and 5 (c) Zero and 4 (d) 3 and 4


Q.31 According to VSEPR theory the shapes of $[\text{SF}_2\text{Cl}]^+$ and $[\text{S}_2\text{O}_4]^{2-}$ should be

- (a) Trigonal planar for $[\text{S}_2\text{O}_4]^{2-}$ and trigonal pyramidal for $[\text{SF}_2\text{Cl}]^+$.
 (b) Both trigonal planar.
 (c) Trigonal pyramidal for $[\text{S}_2\text{O}_4]^{2-}$ and trigonal planar for $[\text{SF}_2\text{Cl}]^+$.
 (d) Both trigonal pyramidal

Q.32 The product of the reaction between $\text{CH}_3\text{Mn}(\text{CO})_5$ and ^{13}CO is

- (a) $(\text{CH}_3^{13}\text{CO})\text{Mn}(\text{CO})_5$ (b) $(\text{CH}_3\text{CO})\text{Mn}(\text{CO})_4(^{13}\text{CO})$
 (c) $(^{13}\text{CH}_3\text{CO})\text{Mn}(\text{CO})_5$ (d) $\text{CH}_3\text{Mn}(\text{CO})_4$


Q.33 The correct pair of ^1H and ^{31}P NMR spectral patterns for $\text{C}(\text{H})(\text{F})(\text{PCl}_2)_2$ is




Q.34 In the following reaction




the major product [X] is




Q.35 In the following reaction




the major product [X] is



Q.36 In the following reaction



the major product [X] is




Q.37 The most appropriate sequence of reactions for carrying out the following conversion is




- (a) (i) Peracid; (ii) H^+ ; (iii) $\text{Zn}/\text{dil. HCl}$.
 (b) (i) Alkaline KMnO_4 ; (ii) NaIO_4 ; (iii) $\text{N}_2\text{H}_4/\text{KOH}$.
 (c) (i) Alkaline KMnO_4 ; (ii) H^+ ; (iii) $\text{Zn}/\text{dil. HCl}$.
 (d) (i) $\text{O}_3/\text{Me}_2\text{S}$; (ii) NaOEt ; (iii) $\text{N}_2\text{H}_4/\text{KOH}$.


Q.38 In the following reaction sequence




the major product [X] is



Q.39 In the following conversion




the major product [X] is



Q.40 In the following reaction




the major product [X] is



Q.41 In the reaction

Optically pure (+)-trans-2-acetoxycyclohexyl tosylate



the major product [X]

- (a) Racemic trans-1,2-cyclohexanediol diacetate.
 (b) Optically active trans-1,2-cyclohexanediol diacetate.
 (c) Racemic cis-1,2-cyclohexanediol diacetate.
 (d) Optically active cis-1,2-cyclohexanediol diacetate.

Q.42 The activity of water at 11 bar and 298 K is

- (a) 1.101 (b) 1.007 (c) 0.998 (d) 0.898

Q.43 For the process



assuming ideal gas behavior, the change in molar entropy is

- (a) $-27.57 \text{ J K}^{-1} \text{ mol}^{-1}$ (b) $+27.57 \text{ J K}^{-1} \text{ mol}^{-1}$ (c) $-24.20 \text{ J K}^{-1} \text{ mol}^{-1}$ (d) $+24.20 \text{ J K}^{-1} \text{ mol}^{-1}$

Q.44 The wave function for a quantum mechanical particle in a 1-dimensional box of length 'a' is given by $\Psi = A \sin \frac{\pi x}{a}$.

The value of 'A' for a box of length 200 nm is

- (a) $4 \times 10^4 \text{ (nm)}^2$ (b) $10\sqrt{2} \text{ (nm)}^{1/2}$ (c) $\sqrt{2}/10 \text{ (nm)}^{-1/2}$ (d) $0.1 \text{ (nm)}^{-1/2}$


Q.45 For 1 mole of a monoatomic ideal gas, the relation between pressure (p), volume (V) and average molecular kinetic energy ($\bar{\epsilon}$) is

- (a) $p = \frac{N_A \bar{\epsilon}}{V}$ (b) $p = \frac{N_A \bar{\epsilon}}{3V}$ (c) $p = \frac{2N_A \bar{\epsilon}}{3V}$ (d) $p = \frac{2N_A \bar{\epsilon}}{3V \bar{\epsilon}}$

Q.46 For a 1 molal aqueous NaCl solution, the mean ionic activity coefficient (γ_{\pm}) and the Debye-Huckel Limiting Law constant (A) are related as

- (a) $\log \gamma_{\pm} = \sqrt{2}A$ (b) $\log \gamma_{\pm} = -\sqrt{2}A$ (c) $\gamma_{\pm} = 10^A$ (d) $\gamma_{\pm} = 10^{-A}$

Q.47 For the concentration cell



the EMF (E) of the cell at a temperature (T) equals

- (a) $2.303 \frac{RT}{F}$ (b) $-2.303 \frac{RT}{F}$ (c) $E_{M^+|M}^0 + 2.303 \frac{RT}{F}$ (d) $E_{M^+|M}^0 - 2.303 \frac{RT}{F}$

Common Data Questions:

Common Data for Questions 48 and 49:

A hypothetical molecule XY has the following properties

Reduced mass: $2 \times 10^{-26} \text{ kg}$

X—Y bond length: 100 pm

Force constant of the bond: $8 \times 10^2 \text{ N.m}^{-1}$

Q.48 The frequency of radiation (in cm^{-1} units) required to vibrationally excite the molecule from $v = 0$ to $v = 1$ state is

- (a) 3184.8 (b) 2123.2 (c) 1061.6 (d) 840.0

Q.49 The frequency of radiation (in cm^{-1} units) required to rotationally excite the molecule from $J = 0$ to $J = 1$ state is

- (a) 1.4 (b) 2.8 (c) 3.2 (d) 3.6

Common Data for Questions 50 and 51:

Na_2HPO_4 and NaH_2PO_4 on heating at high temperature produce a chain sodium pentaphosphate quantitatively.

Q.50 The ideal molar ratio of Na_2HPO_4 to NaH_2PO_4 is

- (a) 4:1 (b) 1:4 (c) 3:2 (d) 2:3




Q.51 The total charge on pentaphosphate anion is

- (a) -5 (b) -3 (c) -7 (d) -9

Linked Answer Questions

Statement for Linked Answer Questions 52 and 53:

The decomposition of ozone to oxygen $2\text{O}_3(\text{g}) \rightarrow 3\text{O}_2(\text{g})$ occurs by the mechanism



where, M is the catalyst molecule. k_i 's are rate constants and $E_{a,i}$'s the activation energies for the elementary steps.

Q.52 Under the steady state approximation for the intermediates, the rate of decomposition of ozone, $-\frac{d[\text{O}_3]}{dt}$, is

- (a) $\frac{2k_1k_3[\text{O}_3]^2[\text{M}]}{k_2[\text{O}_2][\text{M}] + k_3[\text{O}_3]}$ (b) $\frac{2k_1k_3[\text{O}_3]^2[\text{M}]}{k_2[\text{O}_2][\text{M}] - k_3[\text{O}_3]}$
- (c) $\frac{2k_2k_3[\text{O}_3]^2[\text{M}]}{k_2[\text{O}_2][\text{M}] + k_3[\text{O}_3]}$ (d) $\frac{2k_1k_2[\text{O}_3]^2[\text{M}]}{k_2[\text{O}_2][\text{M}] - k_3[\text{O}_3]}$

Q.53 Assuming $k_3[\text{O}_3] \gg k_2[\text{O}_2][\text{M}]$, the activation energy of the overall reaction is


- (a) $\frac{E_{a,1}E_{a,3}}{E_{a,2}}$ (b) $E_{a,3} + E_{a,1} - E_{a,2}$ (c) $E_{a,2}$ (d) $E_{a,1}$

Statement for Linked Answer Questions 54 and 55:


A ketone on treatment with bromine in methanol gives the corresponding monobromo compound [X]

having molecular formula $\text{C}_5\text{H}_9\text{BrO}$. The compound [X] when treated with NaOMe in MeOH produces [Y] as the major product. The spectral data for compound [X] are: ^1H NMR: δ 1.17 (d, 6H), 3.02 (m, 1H), 4.10 (s, 2H); ^{13}C NMR: δ 17, 37, 39, 210.

Q.54 The compound [X] is



Q.55 The major product [Y] is



Section-B

Q.56 – Q.60 carry one mark each.

Q.56 The question below consists of a pair of related words followed by four pairs of words. Select the pair that best expresses the relation in the original pair:

Gladiator : Arena

- (a) Dancer : stage (b) Commuter : train (c) Teacher : classroom (d) Lawyer : courtroom

Q.57 Choose the most appropriate word from the options given below to complete the following sentence:

Under ethical guidelines recently adopted by the Indian Medical Association, human genes are to be manipulated only to correct diseases for which _____ treatments are unsatisfactory

- (a) Similar (b) Most (c) Uncommon (d) Available

Q.58 Choose the word from the options given below that is most nearly opposite in meaning to the given word "Frequency".

- (a) Periodicity (b) Rarity (c) Gradualness (d) Persistency

Q.59 Choose the most appropriate word from the options given below to complete the following sentence:

It was her view that the country's problems had been _____ by foreign technocrats, so that to invite them to come back would be counter-productive.

- (a) Identified (b) Ascertained (c) Exacerbated (d) Analyzed

Q.60 There are two candidates P and Q in an election. During the campaign, 40% of the voters promised to vote for P, and rest for Q. However, on the day of election 15% of the voters went back on their promise to vote for P and instead voted for Q. 25% of the voters went back on their promise to vote for Q and instead voted for P. Suppose, P lost by 2 votes, then what was the total number of voters?

- (a) 100 (b) 110 (c) 90 (d) 95

Q.61 to Q.65 carry two marks each.

Q.61 The horse has played a little known but very important role in the field of medicine. Horses were injected with toxins of diseases until their blood built up immunities. Then a serum was made from their blood. Serums to fight with diphtheria and tetanus were developed this way.

It can be inferred from the passage, that horses were

- (a) Given immunity to diseases (b) Generally quite immune to diseases
(c) Given medicines to fight toxins (d) Given diphtheria and tetanus serums

Q.62 The sum of n terms of the series $4+44+444+\dots$ is

- (a) $(4/81) [10^{n+1} - 9n - 1]$ (b) $(4/81) [10^{n-1} - 9n - 1]$
(c) $(4/81) [10^{n+1} - 9n - 10]$ (d) $(4/81) [10^n - 9n - 10]$


Q.63 Given that $f(y) = |y|/y$, and q is any non-zero real number, the value of $|f(q) - f(-q)|$ is

- (a) 0 (b) -1 (c) 1 (d) 2

Q.64 Three friends, R, S and T shared toffee from a bowl. R took $1/3^{\text{rd}}$ of the toffees, but returned four to the bowl. S took $1/4^{\text{th}}$ of what was left but returned three toffees to the bowl. T took half of the remainder but returned two back into the bowl. If the bowl had 17 toffees left, how many toffees were originally there in the bowl?

- (a) 383 (b) 31 (c) 48 (d) 41

Q.65 The fuel consumed by a motorcycle during a journey while traveling at various speeds is indicated



in the graph below

The distances covered during four laps of the journey are listed in the table below

Lap	Distance (kilometer)	Average speed (kilometer per hour)
P	12	15
Q	75	45
R	40	75
S	10	10

From the given data, we can conclude that the fuel consumed per kilometer was least during the lap

- (a) P (b) Q (c) R (d) S

LEGAL NOTICE

This document is an excerpt from the book entitled “IIT-GATE Chemistry Solved Papers”, and is the intellectual property of the Publisher. The content of this document is protected by international copyright law and is valid only for the personal preview of the user who has originally downloaded it from the publisher’s website (www.dalalinstitute.com). Any act of copying (including plagiarizing its language) or sharing this document will result in severe civil and criminal prosecution to the maximum extent possible under law.



This is a low resolution version only for preview purpose. If you want to read the full book, please consider buying.

Buy the complete book with TOC navigation, high resolution images and no watermark.

Home

CLASSES

NET-JRF, IIT-GATE, M.Sc Entrance & IIT-JAM

Want to study chemistry for CSIR UGC - NET JRF, IIT-GATE, M.Sc Entrance, IIT-JAM, UPSC, ISRO, IISc, TIFR, DRDO, BARC, JEST, GRE, Ph.D Entrance or any other competitive examination where chemistry is a paper ?

[READ MORE](#)

BOOKS

Publications

Are you interested in books (Print and Ebook) published by Dalal Institute ?

[READ MORE](#)

VIDEOS

Video Lectures

Want video lectures in chemistry for CSIR UGC - NET JRF, IIT-GATE, M.Sc Entrance, IIT-JAM, UPSC, ISRO, IISc, TIFR, DRDO, BARC, JEST, GRE, Ph.D Entrance or any other competitive examination where chemistry is a paper ?

[READ MORE](#)

Home: <https://www.dalalinstitute.com/>

Classes: <https://www.dalalinstitute.com/classes/>

Books: <https://www.dalalinstitute.com/books/>

Videos: <https://www.dalalinstitute.com/videos/>

Location: <https://www.dalalinstitute.com/location/>

Contact Us: <https://www.dalalinstitute.com/contact-us/>

About Us: <https://www.dalalinstitute.com/about-us/>

Postgraduate Level Classes (NET-JRF & IIT-GATE)

Admission

[Regular Program](#)

[Distance Learning](#)

[Test Series](#)

[Result](#)

Undergraduate Level Classes (M.Sc Entrance & IIT-JAM)

Admission

[Regular Program](#)

[Distance Learning](#)

[Test Series](#)

[Result](#)

IIT-GATE Chemistry Solved Papers

“IIT-GATE Chemistry Solved Papers” is now available, visit our website for more info.

[READ MORE](#)

Join the revolution by becoming a part of our community and get all of the member benefits like downloading any PDF document for your personal preview.

[Sign Up](#)

Dalal Institute's

IIT-GATE

Chemistry

Solved Papers



DALAL
INSTITUTE

2011-2020 & Letest Model Test


Table of Contents

IIT-GATE: Model Test	5
Chemistry	5
❖ Question Paper.....	5
❖ Answer Key	19
❖ Solution.....	20
IIT-GATE: 2011	24
Chemistry	24
❖ Question Paper.....	24
❖ Answer Key	41
❖ Solution.....	42
IIT-GATE: 2012	46
Chemistry	46
❖ Question Paper.....	46
❖ Answer Key	63
❖ Solution.....	64
IIT-GATE: 2013	68
Chemistry	68
❖ Question Paper.....	68
❖ Answer Key	83
❖ Solution.....	84
IIT-GATE: 2014	90
Chemistry	90
❖ Question Paper.....	90
❖ Answer Key	106
❖ Solution.....	107
IIT-GATE: 2015	111
Chemistry	111
❖ Question Paper.....	111

❖ Answer Key	126
❖ Solution.....	127
IIT-GATE: 2016	131
Chemistry	131
❖ Question Paper.....	131
❖ Answer Key	150
❖ Solution.....	151
IIT-GATE: 2017	155
Chemistry	155
❖ Question Paper.....	155
❖ Answer Key	171
❖ Solution.....	172
IIT-GATE: 2018	175
Chemistry	175
❖ Question Paper.....	175
❖ Answer Key	191
❖ Solution.....	192
IIT-GATE: 2019	196
Chemistry	196
❖ Question Paper.....	196
❖ Answer Key	216
❖ Solution.....	217
IIT GATE: 2020.....	222
Chemistry	222
❖ Question Paper.....	222
❖ Answer Key	239
❖ Solution.....	240

The best institute for CSIR-JRF, UGC-NET, IIT-GATE, IIT-JAM, UPSC, GRE, IISc, TIFR, DRDO, BARC, JEST, ISRO and all Ph.D-M.Sc entrance examinations where chemistry is a paper.

*Dalal Institute's
other publications in this series*



D DALAL
INSTITUTE

Main Market, Sector 14, Rohtak, Haryana 124001, India
(info@dalalinstitute.com, +91-9802825820)
www.dalalinstitute.com